

Gin Tonic

Reference: FD 18 03C

INCI name	%	Function
PHASE A		
Water (<i>Aqua</i>)	up to 100.00	
Glycerin	3.00	Humectant
Sodium Polyacrylate	0.40	Thickener
Tapioca Starch	0.20	Thickener
PHASE B		
Polyglyceryl-2 Stearate (<i>and</i>) Glyceryl Stearate (<i>and</i>) Stearyl Alcohol - PolyAqual™ 2W	3.00	Emulsifier
Isodecyl Neopentanoate	3.00	Emollient
Propylheptyl Caprylate	4.00	Emollient
Octyldodecyl Oleate (<i>and</i>) Octyldodecyl Stearoyl Stearate (<i>and</i>) Polyhydroxystearic Acid (<i>and</i>) Octyldodecanol - innollient™ LO	3.00	Emollient
Butyrospermum Parkii (<i>Shea</i>) Butter	1.00	Emollient
PHASE C		
Dimethicone	1.00	Emollient
PHASE D		
Glycerin (<i>and</i>) Water (<i>and</i>) Panax Ginseng Root Extract - GinsenoLite™-G	2.00	Active Ingredient
PHASE E		
Preservative	a.n.	Preservative
Parfum	a.n.	Parfum
VISCOSITY after 24h Brk, RVDV-I Prime, Sp 93, 10 rpm 20000 – 30000 cps		pH (100%) 5.0 – 7.0

Procedure:

- Prepare phase A and phase B separately and warm them up to 75°C;
- Disperse the thickener in the water phase using a suitable dispersion unit;
- Add phase B to phase A and homogenize for a few minutes;
- Add phase C and homogenize for a few minutes;
- Cool down to 40°C under gentle stirring and add phases D and E.